

Fluctuation Effects of Gauge Fields in the Slave-Boson t-J Model

Ikuo Ichinose^{*,1} Tetsuo Matsui^{†,2} and Kazuhiko Sakakibara^{*3}

^{*}Institute of Physics, University of Tokyo, Komaba, Tokyo, 153 Japan

[†]Department of Physics, Kinki University, Higashi-Osaka, 577 Japan

^{*}Department of Physics, Nara National College of Technology, Yamatokohriyama,
639-11 Japan

Abstract

We present a quantitative study of the charge-spin separation (CSS) phenomenon in a U(1) gauge theory of the t-J model of high- T_C superconductors. We calculate the critical temperature T_{CSS} of a confinement-deconfinement phase transition as a function of the hole doping δ , below which the CSS takes place. The fluctuations of gauge field are so large that T_{CSS} is reduced to about 10 % of its mean-field value.

¹e-mail address: ikuo@hep1.c.u-tokyo.ac.jp

²e-mail address: matsui@phys.kindai.ac.jp

³e-mail address: sakaki@center.nara-k.ac.jp

The anomalous properties of the metallic phase of cuprate superconductors, like the linear temperature (T) dependence of resistivity, have called for theoretical explanations. The idea of charge-spin separation (CSS) of electrons into holons and spinons, introduced by Anderson[1], seems to be an interesting possibility for them.

Straightforward mean-field (MF) theories of the t-J model in the slave-boson (SB) or slave-fermion representation are compatible with the CSS since holons and spinons acquire their own hopping amplitudes through nonvanishing MF's. Fluctuations around such MF's are described by a U(1) gauge theory. Effects of these fluctuations have been studied within perturbative approaches. In Ref.[2] the resistivity is calculated to show a linear- T dependence. From a more general point of view, a system of nonrelativistic fermions coupled with a U(1) gauge field is studied by perturbative renormalization-group (RG), showing the non-Fermi-liquid-like behavior[3]. All these analyses assume that gauge-field fluctuations are “*small*” and can be handled by perturbative or RG methods.

In Ref.[4], we developed a general formalism to study such gauge fields in a *non-perturbative* manner, in order to see whether and when these perturbative analyses are validated. In Ref.[5], we applied that formalism to the t-J model. Our method was motivated by the success of the Ginzburg-Landau (GL) theory of the microscopic Bardeen-Cooper-Schrieffer's (BCS) model, which, taking a form of the XY spin model, is useful for studying superconducting phase transition, dynamics of vortices, etc.

In contrast to the XY model for the BCS model, the resulting GL theory for the t-J model takes a form of lattice gauge theory, reflecting the local gauge symmetry in the slave-particle representation, so the gauge-theoretical approach is quite useful[6]. Gauge dynamics generally exhibits two phases; confinement phase with large fluctuations of gauge fields and deconfinement phase with small fluctuations. In the language of gauge theory, we characterized CSS as a deconfinement phenomenon of holons and spinons[4]. By using the knowledge of lattice gauge theory, we showed that there exists a confinement-deconfinement (CD) phase transition at some critical temperature

T_{CSS} . At $T < T_{\text{CSS}}$, the deconfinement phase is realized, where holons and spinons are deconfined, hence the CSS takes place and the anomalous behavior is expected, while at $T > T_{\text{CSS}}$, the confinement phase is realized where holons and spinons are confined in electrons and the usual Fermi-liquid picture of electrons should be applicable.

In this paper, we calculate T_{CSS} quantitatively following Ref.[5]. The result should be compared with the experimental indications of the onset T , below which the anomalous behavior is observed.

In the SB representation of the t-J model, the electron operator $C_{x\sigma}$ [x ; site, $\sigma(= 1, 2)$; spin] is written as $C_{x\sigma} = b_x^\dagger f_{x\sigma}$, where $f_{x\sigma}$ are the fermionic spinon operators and b_x are the bosonic holon operators. The physical states must satisfy the local constraint $(f_{x1}^\dagger f_{x1} + f_{x2}^\dagger f_{x2} + b_x^\dagger b_x - 1)|\text{phys}\rangle = 0$ to avoid double occupancies of electrons. The Hamiltonian on a two-dimensional lattice is given by

$$\begin{aligned} H &= -t \sum_{x,i,\sigma} (b_{x+i}^\dagger f_{x\sigma}^\dagger f_{x+i,\sigma} b_x + \text{H.c.}) \\ &- \frac{J}{2} \sum_x \left(\sum_\sigma f_{x\sigma}^\dagger \tilde{f}_{x+i,\sigma} \right) \left(\sum_{\sigma'} \tilde{f}_{x+i,\sigma'}^\dagger f_{x\sigma'} \right) + H_\mu, \end{aligned} \quad (1)$$

where $i(= 1, 2)$ is the direction index, $\tilde{f}_{x\sigma} \equiv \sum_{\sigma'} \epsilon_{\sigma\sigma'} f_{x\sigma'}^\dagger$ ($\epsilon_{12} = -\epsilon_{21} = 1$), and $H_\mu = -\mu_f \sum f_{x\sigma}^\dagger f_{x\sigma} - \mu_b \sum b_x^\dagger b_x$. We relaxed the local constraint to the global one. The chemical potentials μ_b and μ_f are chosen as $\langle b_x^\dagger b_x \rangle = \delta$ and $\langle \sum_\sigma f_{x\sigma}^\dagger f_{x\sigma} \rangle = 1 - \delta$.

By using the Hubbard-Stratonovich transformation in the path-integral formalism, we rewrite (1) in the form of MF theory proposed in Ref.[7]. Explicitly, by introducing auxiliary complex *link variables*, χ_{xi} and D_{xi} which are *fluctuating* “MF’s”, we get

$$\begin{aligned} H &= \sum_{x,i} \left[\frac{3J}{8} |\chi_{xi}|^2 + \frac{2}{3J} |D_{xi}|^2 \right] \\ &- \sum_{x,i} \left[\chi_{xi} \left(\frac{3J}{8} \sum_\sigma f_{x+i,\sigma}^\dagger f_{x\sigma} + t b_{x+i}^\dagger b_x \right) + \text{H.c.} \right] \\ &- \frac{1}{2} \sum_{x,i,\sigma} [D_{xi} f_{x\sigma}^\dagger \tilde{f}_{x+i,\sigma} + \text{H.c.}] + H_4 + H_\mu, \end{aligned} \quad (2)$$

where $H_4 = \frac{8t^2}{3J} \sum b_{x+i}^\dagger b_{x+i} b_x^\dagger b_x - \frac{3J}{8} \sum f_{x+i,\sigma}^\dagger f_{x+i,\sigma'} f_{x\sigma'}^\dagger f_{x\sigma}$. The partition function

$Z(\beta)$ [$\beta \equiv (k_B T)^{-1}$] is given by

$$\begin{aligned} Z &= \int [db][df][d\chi][dD] \exp(A), \\ A &= \int_0^\beta d\tau [-\sum_x (\bar{b}_x \dot{b}_x + \sum_\sigma \bar{f}_{x\sigma} \dot{f}_{x\sigma}) - H]. \end{aligned} \quad (3)$$

From (3), one gets the relations, $\langle \bar{\chi}_{xi} \rangle = \langle \sum f_{x+i,\sigma}^\dagger f_{x\sigma} + (8t/3J) b_{x+i}^\dagger b_x \rangle$, $\langle \bar{D}_{xi} \rangle = (3J/4) \langle \sum f_{x\sigma}^\dagger \tilde{f}_{x+i,\sigma} \rangle$. So χ_{xi} is the hopping amplitude of holons and spinons, while D_{xi} is the amplitude of resonating valence bonds (RVB) of antiferromagnetism.

In Ref.[5], we derived an equation that determines T_{CSS} . Here, we summarize the basic steps (I-IV) leading to that equation.

(I) We parametrize χ_{xi} and D_{xi} a la “uniform RVB state” as $\chi_{xi} = \chi U_{xi}$, $D_{xi} = D_i V_{xi}$, where $U_{xi}, V_{xi} \in U(1)$, and $\chi, D_1 = -D_2 = D$ are link-independent constants, focusing on the U(1) phase dynamics. Eq.(2) is invariant under a time-independent U(1) gauge transformation, $(f_{x\sigma}, b_x) \rightarrow e^{i\theta_x}(f_{x\sigma}, b_x)$, while the above U(1) variables transform as $U_{xi} \rightarrow e^{i\theta_x} U_{xi} e^{-i\theta_{x+i}}$ and $V_{xi} \rightarrow e^{i\theta_x} V_{xi} e^{i\theta_{x+i}}$. Therefore they are two different kinds of lattice gauge variables.

(II) We integrate over $f_{x\sigma}$ and b_x by using the hopping expansion, an expansion in powers of χ_{xi} and D_{xi} . By keeping the leading contributions (i.e., neglecting the plaquette terms, etc.), we obtain the following effective lattice gauge theory as the GL theory of the t-J model;

$$\begin{aligned} Z &= \int [dU][dV] \exp(A_{\text{eff}}), \\ A_{\text{eff}} &= \int_0^\beta d\tau \sum_{x,i} [-a_U |\dot{U}_{xi}|^2 - a_V |\dot{V}_{xi}|^2], \end{aligned} \quad (4)$$

where $a_{U,V}$ are known functions of $\chi, D, \mu_{b,f}$ and T .

(III) Following the Polyakov-Susskind approach[8] to the CD transition we map (4) into an *anisotropic* 2D classical XY spin model;

$$\begin{aligned} Z &= \int_{-\pi}^\pi \prod_x \frac{d\alpha_x}{2\pi} \exp[\sum_{x,i} \{J_1 \cos(\alpha_{x+i} - \alpha_x) \\ &\quad + J_2 \sum_{x,i} \cos(\alpha_{x+i} + \alpha_x)\}], \end{aligned} \quad (5)$$

where $J_1 \equiv \beta^{-1}a_U$, $J_2 \equiv \beta^{-1}a_V$. It is obvious that $J_2 \propto D^2$ is an anisotropic parameter. The spin-spin correlation $C(|x|) \equiv \langle \exp(i\alpha_x) \exp(-i\alpha_0) \rangle$ is expressed as $\exp(-\beta W(|x|))$, where $W(R)$ is the potential energy of a pair of oppositely charged gauge sources separated by R .

(IV) We investigated the spin model (5) both by the effective action in large-spatial-dimensions [5] and the Monte Carlo simulation[9]. From both analyses, we found that there occurs an order-disorder phase transition along the line $J_1 + J_2 \simeq 1$, which is identified with the CD phase transition. The condition $J_1 + J_2 \simeq 1$ gives rise to the following equation that determines T_{CSS} [10];

$$(k_B T_{\text{CSS}})^2 \simeq \frac{\chi^2(T_{\text{CSS}})}{4\pi^2} \left[\frac{1}{2} \left(\frac{3J}{8} \right)^2 (1 - \delta^2) + t^2 \delta (1 + \delta) \right] + \frac{D^2(T_{\text{CSS}})}{32F^3} [2\delta - (1 - \delta)^2 F(F + 1)], \quad (6)$$

where $F \equiv \ln((1 + \delta)/(1 - \delta))$. There are three phases of the spin dynamics (5) and corresponding gauge dynamics[11], which are summarized as follows.

- Disordered Phase at $J_1 + J_2 < 1$:

$C(R) \sim \exp(-\gamma R)$ with some finite constant γ , and so $W(R) \propto R$ is a confining potential. The gauge system is in the *confinement phase* ($\langle U_{xi} \rangle = \langle V_{xi} \rangle = 0$), where only the gauge-invariant neutral objects, like electrons, appear.

- Ordered Phase at $J_1 + J_2 > 1$, $J_2 \neq 0$:

The Z_2 symmetry of (5) is spontaneously broken. Excitations from the ground state $\alpha_x = 0 \pmod{\pi}$ are *massive* spin waves with finite energy gap m ; $C(R) \rightarrow \text{const.} + \exp(-mR)$. Then, $W(x) \simeq \exp(-m|x|)$. This is the *deconfinement* ($\langle U_{xi} \rangle, \langle V_{xi} \rangle \neq 0$) *Higgs phase* where the gauge fields acquire a mass m .

- Quasi-Ordered Phase at $J_1 > 1$, $J_2 = 0$:

The model is $O(2)$ symmetric and the Kosterlitz-Thouless analysis applies; $C(R) \simeq R^{-\eta}$, $W(R) \simeq \ln(R)$. This is the *deconfinement Coulomb phase* with massless gauge fields.

To determine T_{CSS} , we first calculate χ , D , $\mu_{b,f}$ as functions of T and δ by a

straightforward MF theory that totally ignores gauge-field fluctuations [12]. In Figs. 1 and 2, we summarize the results of such a MF theory with $J = 0.1eV, t = 0.3eV$. Fig.1 shows $J_1 + J_2$ as a function of T for various δ 's. In Fig.2, the curves T_χ and T_{RVB} show the value of T at which χ and D vanish, respectively. T_{BC} is the onset T , below which the Bose condensation $\langle b_x \rangle \neq 0$ takes place. To calculate T_{BC} we included a weak but finite three-dimensionality, $t' = \alpha t, \alpha \simeq 10^{-2}$, while χ, D are calculated with $\alpha = 0$. Using these χ, D , we can solve (6) to obtain T_{CSS} as a function of δ as plotted in Fig.2. We find that T_{CSS} is smaller by factor $\sim 1/10$ than its mean-field value T_χ . Fig.3 shows this relation together with $\chi(T)$. Figs.2,3 exhibit that the effects of gauge fluctuations on T_{CSS} are so strong that the MF estimation is quantitatively incorrect. This result is welcome since the experimental signal of the onset of CSS is around $T \sim$ several-hundred K .

To check the stability of T_{CSS} , let us suppose that the condition for the transition would be $J_1 + J_2 = A$ with $A = 2$ instead of $A = 1$, due to, say, the higher-order terms in the hopping expansion. Fig.1 shows that T_{CSS} would shift only $20 \sim 30\%$ from its present value since $J_1 + J_2$ decreases very rapidly up to $J_1 + J_2 \simeq 0.5$. This reflects the fact that $\chi(T)$ is almost saturated in this *low* T region, as shown in Fig.3.

The two-body term H_4 in (2), neglected in the MF theory, is necessary to keep the equivalence to the original t-J model. In the leading-order of perturbation theory, it reduces T_{CSS} furthermore. Also one may ask how T_{CSS} changes in other decoupling schemes [13]. Although T_{CSS} may certainly shift, the importance of large fluctuations should remain true. We shall report on these points in future publications.

The concept of CSS is very drastic, and it is difficult to handle it by the conventional approaches in the condensed matter physics. However, the notion of gauge theory affords us to understand it as a familiar phenomenon, like the CD phenomenon of quarks and gluons in the color gauge theory of strong interactions. We note that the gauge theory of separation phenomena used here found yet another application; the particle-flux separation in the fractional quantum Hall effects[14].

References

- [1] P.W.Anderson, Phys.Rev.Lett.**64**, 1839(1990).
- [2] P.Lee and N.Nagaosa, Phys.Rev.**B46**,5621(1992).
- [3] C.Nayak and F.Wilczek, Nucl.Phys.**B417**[FS],359 (1994);
ibid. **B430**[FS], 534(1994);
S.Chakravarty, R.E.Norton and O.F.Syljuasen, Phys.Rev.Lett.**74**, 1423(1995);
I.Ichinose and T.Matsui, Nucl.Phys.**B441**[FS], 483(1995);
M.Onoda, I.Ichinose and T.Matsui, Nucl.Phys.**B446**[FS], 353(1995).
- [4] I.Ichinose and T.Matsui, Nucl.Phys.**B394**,281 (1993).
- [5] I.Ichinose and T.Matsui, Phys.Rev.**B51**,11860(1995).
- [6] The GL theory for superconductivity was discussed by G.Baskaran and P.W.Anderson, Phys.Rev. **B37**, 580(1988), A.Nakamura and T.Matsui, Phys.Rev. **B37**,7940(1988), I.Ichinose and T.Matsui, Phys.Rev. **B45**,9976(1992). The implication of gauge symmetry in the normal channel was discussed in L.B.Ioffe and I.Larkin, Phys.Rev. **B39**,8988(1989). The GL theory of the present paper is for the CSS transition.
- [7] M.U.Ubbens and P.A.Lee, Phys.Rev.**B46**, 8434(1992); *ibid.* **B49**, 6853(1994).
- [8] A.M.Polyakov, Phys.Lett.**B72**,477(1978);
L.Susskind, Phys.Rev. **D20**,2610(1979).
- [9] K.Sakakibara and T.Matsui, unpublished.
- [10] The corresponding Eq.(3.30) in Ref.[5] should be corrected as (6). This reflects the errata; The L.H.S. of the first line in Eq.(3.8) should be multiplied by a factor 2. Also the minus signs of $\mu_{f,b}$ in the Hamiltonian (3.2,3) should be reversed. To

obtain (6), we used $\mu_{b,f}(\delta)$ determined by the hopping expansion, not by the MFT.

- [11] One might conceive the possibility of mixed dynamics; the U field (say, $b - \bar{b}$ channel) is in the confinement phase whereas the V ($b - b$ channel) field is in the deconfinement (or otherway around). However, a MF theory and strong-coupling expansion for (5) shows $C(|x|)$ and $\langle \exp(i\alpha_x) \exp(i(-)^{x_1+x_2}\alpha_0) \rangle$ behave similarly, denying such a possibility.
- [12] Our MF theory neglected H_4 of (2) as in [7]. We assumed the uniform RVB, without setting the spatial phases as variational parameters as in their case.
- [13] Kotliar and Liu, Phys.Rev. **B38**,5142(1988);
Y.Suzumura, Y.Hasegawa, and H.Fukuyama, J.Phys.Soc.Jpn.**57**, 2768(1988).
- [14] I.Ichinose and T.Matsui, Nucl.Phys. **B468**[FS],487(1996);
*ibid.***B483**[FS],681(1997).

FIGURE CAPTIONS.

Fig.1. $J_1 + J_2$ v.s. T . The CSS sets in at $J_1 + J_2 \simeq 1$.

Fig.2. MF results and T_{CSS} . Along T_χ and T_{RVB} , χ and D vanish, respectively. T_{BC} is the onset T of Bose condensation, $\langle b_x \rangle \neq 0$. T_{CSS} is the critical T of the CD transition calculated by Eq.(6), *below* which the CSS takes place. The dashed line starting from P is an *expected* T along which $\langle D_{xi} \rangle$ vanishes, i.e., T_{RVB} renormalized by gauge fluctuations $\langle V_{xi} \rangle$. Similarly, the renormalized T_χ is just T_{CSS} . There are five phases:

- (i) Strange Metal Phase: Deconfinement-Coulomb;
- (ii) Spin-Gap Phase: Deconfinement-Higgs with $D \neq 0$;
- (iii) Fermi-Liquid Phase: Deconfinement-Higgs with $\langle b_x \rangle \neq 0$;
- (iv) Superconducting Phase: Deconfinement-Higgs with $D, \langle b_x \rangle \neq 0$;
- (v) Electron Phase: Confinement *above* T_{CSS} .

Fig.3. $\chi(T)$ at $\delta = 0.15$. The overall structure of $\chi(T)$ and the relation $T_{\text{CSS}} \sim T_\chi/10$ remain similar for other values of $0 < \delta < 1$.



